

## Design and Synthesis of New Barbituric- and Thiobarbituric Acid Derivatives as Potent Urease Inhibitors: Structure Activity Relationship And Molecular Modeling Studies

Abdul Rauf<sup>a</sup>, Sohail Shahzad<sup>a,b</sup>, Marek Bajda<sup>c</sup>, Muhammad Yar<sup>b</sup>, Faiz Ahmed<sup>a</sup>, Nazar Hussain<sup>a</sup>, Muhammad Nadeem Akhtar<sup>d</sup>, Ajmal Khan<sup>e</sup>, Jakub Jończyk<sup>c</sup>

<sup>a</sup> Department of Chemistry, The Islamia University of Bahawalpur, Bahawalpur 63100, Pakistan

<sup>b</sup> Interdisciplinary Research Center in Biomedical Materials, COMSATS Institute of Information Technology, Lahore 54000, Pakistan

<sup>c</sup> Department of Physicochemical Drug Analysis, Faculty of Pharmacy, Jagiellonian University Medical College, Medyczna 9, 30-688 Cracow, Poland

<sup>d</sup> Faculty of Industrial Sciences & Technology, Universiti Malaysia Pahang, Lebuhraya Tun Razak 26300, Kuantan, Pahang, Malaysia

<sup>e</sup> H.E.J. Research Institute of Chemistry, International Center for Chemical and Biological Sciences, University of Karachi, Karachi 75270, Pakistan

### ABSTRACT

In this study 36 new compounds were synthesized by condensing barbituric acid or thiobarbituric acid and respective anilines (bearing different substituents) in the presence of triethyl orthoformate in good yields. In vitro urease inhibition studies against jack bean urease revealed that barbituric acid derived compounds (**1–9** and **19–27**) were found to exhibit low to moderate activity however thiobarbituric acid derived compounds (**10–18** and **28–36**) showed significant inhibition activity at low micro-molar concentrations. Among the synthesized compounds, compounds (**15**), (**12**), (**10**), (**36**), (**16**) and (**35**) showed excellent urease inhibition with IC<sub>50</sub> values  $8.53 \pm 0.027$ ,  $8.93 \pm 0.027$ ,  $12.96 \pm 0.13$ ,  $15 \pm 0.098$ ,  $18.9 \pm 0.027$  and  $19.7 \pm 0.63$   $\mu\text{M}$ , respectively, even better than the reference compound thiourea (IC<sub>50</sub> =  $21 \pm 0.011$ ). The compound (**11**) exhibited comparable activity to the standard with IC<sub>50</sub> value  $21.83 \pm 0.19$   $\mu\text{M}$ . In silico molecular docking studies for most active compounds (**10**), (**12**), (**15**), (**16**), (**35**) and (**36**) and two inactive compounds (**3**) and (**6**) were performed to predict the binding patterns.

**KEYWORDS:** Urease; Barbituric; Thiobarbituric; Jack bean; In silico; Triethyl orthoformate

**DOI:** [10.1016/j.bmc.2015.05.038](https://doi.org/10.1016/j.bmc.2015.05.038)